



Factsheet #10



Mapping vegetation with remote sensing

Mapping vegetation types


Mapping vegetation from the air or from space is possible using a growing number of tools. These cover a wide range of spatial and temporal scales, from one-time airplane campaigns to continuous satellite measurements over the past 40 years.

One key requirement to classifying satellite or other remote sensing data is **field verification campaigns**. To properly translate what you can see from afar, you need to have detailed information from the ground.

During the course of the ORYCS project, such field verification data was collected, resulting in a high-resolution map of major vegetation types in a test area in Northern Namibia.



Collecting detailed verification data is essential. Data on plant type, density, height, and health can be used to interpret remotely-sensed data over a larger spatial area. This makes it possible to go from sparse data to a continuous map covering a larger region.




Provisional map of major vegetation types southwest of the Etosha Pan, based on Sentinel-2 data and ~100 field verification points. Coordinates in UTM 33S. Etosha Heights and Etosha National Park boundaries shown for reference.

Etosha Heights Reserve	<i>Catophractes alexandri</i>	Open grassland
Etosha National Park	Mopane medium/large	Small open areas with calc. Rocks
<i>Senegalia (Acacia) mellifera</i>	Mopane open savanna	Mixed stands on plain outcrops
<i>Vachellia (Acacia) newbournii</i>	Mopane riparian	Mixed stands on mountains
<i>Vachellia (Acacia) reficiens</i>	Mopane small	Bare soil

Mapping individual trees

Improvements in **image resolution**, for example from air photos, also allow for new techniques to be applied. When high-resolution data is available, it can be used to map vegetation at an unprecedented scale.



The above aerial image illustrates how **machine learning** methods can be used to map individual tree locations over a large area. These cannot only be used to improve species maps, but also biomass estimations. The extent of this figure is 100×100 m with a spatial resolution of 10×10 cm. Not all trees are perfectly mapped – ongoing adjustments to the training data and mapping algorithm will improve this process with the goal of providing a complete tree accounting alongside precise tree type classifications.


References

Atmani, F., Bookhagen, B., Smith, T. (2022). Measuring Vegetation Heights and Their Seasonal Changes in the Western Namibian Savanna Using Spaceborne Lidars. *Remote Sensing*, 14(12), 2928. <https://doi.org/10.3390/rs14122928>

Hering, R., Hauptfleisch, M., Jago, M., Smith, T., Kramer-Schadt, S., Stiegler, J., Blaum, N. (2022). Don't stop me now: Managed fence gaps could allow migratory ungulates to track dynamic resources and reduce fence related energy loss. *Frontiers in Ecology and Evolution*, 1–18. <https://doi.org/10.3389/fevo.2022.907079>

Vegetation heights and biomass

There is an increasing amount of **Lidar** data (light detection and ranging), collected from the air and from space. These data measure precise **heights** using a laser, and can also be used to infer the **density** of plants.



These measurements were collected from space via laser measurements, and show the precise height of vegetation above the ground. Observations are collected along a profile, hence they are not spatially continuous (as is usual for optical data). Such data can be used to map **biomass** across large areas. This figure shows only vegetation heights above 3m to highlight tree occurrence. Combined with other spaceborne data, these can be used to monitor **annual cycles in vegetation growth** – even for shorter vegetation such as grass and shrubs.

High-resolution photos can also be used to create high accuracy **3D models**, which complement space-based approaches. Combining these vegetation-density estimates with tree-counting or vegetation species maps will allow new and high-resolution studies on large-scale ecology in savannah ecosystems.

Authors

Taylor Smith (University of Potsdam), **Bodo Bookhagen** (University of Potsdam), **Robert Hering** (University of Potsdam), **Tim Herkenrath** (University of Potsdam)

The ORYCS Project

The German-Namibian research project “ORYCS – Options for sustainable land use adaptations in savanna systems” aims to assess the suitability of wildlife management strategies in Namibia as options for adapting land use to climate change in savanna ecosystems.

www.orycs.org

The ORYCS project is part of the SPACES II research program and is funded by the German Federal Ministry of Education and Research (BMBF) under the funding reference FKZ 01LL1804A-C.

Project Coordination

German Coordination

PD Dr. Niels Blaum
E-Mail: blaum@uni-potsdam.de

Namibian Coordination

Dr. Morgan Hauptfleisch
E-Mail: mhauptfleisch@nust.na



Published by:
ORYCS Project Consortium
Frankfurt am Main, Potsdam, Berlin, Windhoek

May 2023



Attribution-NonCommercial 3.0 Germany
(CC BY-NC 3.0 DE)

